

THE q -BOSON REALIZATIONS OF THE QUANTUM GROUP $U_q(sl(n+1, C))$

Č Burdík[◇], L.Černý^{*} and O. Navrátil^{**}

[◇] Nuclear Centre, Faculty of Mathematics and Physics, Charles University,
V Holešovičkách 2, 180 00 Prague 8, Czechoslovakia

^{*} Faculty of Nuclear Sciences and Physical Engineering, Technical University of
Prague, Trojanova 13, 120 00 Prague 2, Czechoslovakia
and

^{**} Institute of Geotechnics, Czechoslovak Academy of Sciences, V Holešovičkách
41, 182 09 Prague 8, Czechoslovakia

Abstract

We give explicit expression of recurrency formulae of canonical realization for quantum enveloping algebras $U_q(sl(n+1, C))$. In these formulas the generators of the algebra $U_q(sl(n+1, C))$ are expressed by means of n -canonical q -boson pairs one auxiliary representation of the algebra $U_q(gl(n, C))$.

In a recent paper (Fu and Ge 1992) the authors gave a general method to construct the q-boson realizations of quantum algebras from their Verma representations. The method was illustrated on two examples of algebras $U_q(sl(2, C))$ and $U_q(sl(3, C))$. In the case of Lie algebras this method was formulated in (Burdík 1985) and some generalization for $U_q(sl(2, C))$ in (Burdík a Navrátil 1990).

In this letter we are devoted to studying explicitly the general case $U_q(sl(n + 1, C))$. Because it is difficult to write down the explicit expression of its Verma representation we use the recurrency from $U_q(gl(n + 1, C))$ to $U_q(gl(n, C))$.

In final formulae the generators of the algebra $U_q(sl(n + 1, C))$ are expressed by means of n-canonical boson pairs and auxiliary representation of the algebra $U_q(gl(n, C))$. The pure q-boson realizations we can obtain than after recurrency.

Very similar formulae were used in our paper (Burdík, Havlíček and Vančura 1992) for construction of irreducible highest weight representations of quantum groups $U_q(gl(n + 1, C))$.

Now it is clear (Fu and Ge 1992) that these formulas we can used for construction of parametrized cyclic representations starting from cyclic representations of q-deformed Weyl algebra. The irreducibility of these representations we study and the results will be published.

We believe that our recurrency method can be used to construct q-boson realizations for deformations of other semi-simple Lie algebras as well, and some positive indication for deformation of B_n and D_n have already been obtained.

The q-Weyl algebras are defined as associative algebras W_2^q over C generated by b^+, b , and $q^{\pm N}$ satisfying (Hayshi 1990)

$$bb^+ - q^{\mp}b^+b = q^{\pm N}, q^N q^{-N} = q^{-N} q^N = 1,$$

$$q^N b^{\pm} q^{-N} = q^{\pm} b^{\pm}, (b^- = b),$$

which degenerates to the usual W algebras in the limit $q \rightarrow 1$.

The n-pairs Weyl algebra we obtain as

$$W_{2n}^q = W_2^q \otimes W_2^q \otimes \dots \otimes W_2^q \quad n - times$$

and the different pairs commute.

The quantum group $U_q(sl(n + 1, C))$ is defined by the generators k_i, k_i^{-1}, e_i , and f_i for $i = 1, \dots, n$, and the relations

$$k_i k_i^{-1} = k_i^{-1} k_i = 1, \quad k_i k_j = k_j k_i,$$

$$k_i e_j k_i^{-1} = q^{a_{ij}} e_j, \quad k_i f_j k_i^{-1} = q^{-a_{ij}} f_j,$$

$$[e_i, f_j] = \delta_{ij} \frac{k_i - k_i^{-1}}{q - q^{-1}}, \quad (1)$$

$$[e_i, e_j] = [f_i, f_j] = 0 \text{ for } |i - j| \geq 2,$$

$$e_i^2 e_{i\pm 1} - (q + q^{-1}) e_i e_{i\pm 1} e_i + e_{i\pm 1} e_i^2 = 0,$$

$$f_i^2 f_{i\pm 1} - (q + q^{-1}) f_i f_{i\pm 1} f_i + f_{i\pm 1} f_i^2 = 0,$$

where $(a_{ij})_{i,j=1,\dots,n}$ is the Cartan matrix of $U_q(sl(n+1, C))$, i.e. $a_{ii} = 2, a_{i\pm 1, i} = -1$ and $a_{ij} = 0$ for $|i - j| \geq 2$.

The generators e_i, f_i correspond to the simple roots. According to (Rosso 1988) and (Burroughs 1990) we introduce the generators

$$X_n = e_n$$

and recurrently

$$X_r = e_r X_{r+1} - q X_{r+1} e_r \text{ for } r = 1, \dots, n-1. \quad (2)$$

The generators k_i, k_i^{-1}, e_i, f_i for $i = 1, 2, \dots, n-1$ and k_n form a subalgebra in $U_q(sl(n+1, C))$ and evidently it is $U_q(gl(n, C))$. We add to this subalgebra the generator f_n and we obtain again the subalgebra which we will denote A .

There exists a very simply representation ϕ of A in $U_q(gl(n, C))$

$$\phi(z)y = z.y \quad \phi(f_n)y = 0 \text{ for any } z, y \in U_q(gl(n, C))$$

It is a left regular representation. Because

$U_q(sl(n+1, C))(A \otimes z - 1 \otimes \phi(A)z) \quad z \in U_q(gl(n, C))$ is an invariant subspace of the left regular of $U_q(sl(n+1, C))$ we can define a generalized Verma representation ϱ as a factor representation of the left regular of $U_q(sl(n+1, C))$ with respect to this subspace.

The representation space of the representation ϱ is given by

$$V(\lambda_n) = (X_n)^{m_n} (X_{n-1})^{m_{n-1}} \dots (X_1)^{m_1} \otimes U_q(gl(n, C))$$

We will denote

$$|m\rangle \otimes w = |m_n, m_{n-1}, \dots, m_1\rangle \otimes w = (X_n)^{m_n} (X_{n-1})^{m_{n-1}} \dots (X_1)^{m_1} \otimes w$$

where $w \in U_q(gl(n, C))$ and define the representation Γ of $W_{2(n)}$ on $V(\lambda_n)$ by

$$\Gamma(b_i^+) |m\rangle \otimes w = |m+1_i\rangle \otimes w = |m_n, \dots, m_i+1, \dots, m_1\rangle \otimes w,$$

$$\Gamma(b_i) |m\rangle \otimes w = [m_i]_q |m-1_i\rangle \otimes w = [m_i]_q |m_n, \dots, m_i-1, \dots, m_1\rangle \otimes w, \quad (3)$$

$$\Gamma(q^{N_i}) |m\rangle \otimes w = q^{m_i} |m\rangle \otimes w,$$

$$\Gamma([N_i + \alpha])|m\rangle \otimes w = [m_i + \alpha] |m\rangle \otimes w.$$

Now for explicit construction we will need the commutation relations X_j with e_i, f_i and k_i . Starting from here we will take $r < n$.

Evidently

$$e_r X_s = X_s e_r \text{ for } r < s - 1 \quad (4)$$

because in X_s are included only e_t for $t > s + 1$ which commute with e_r . For further calculation the following lemma will be useful.

Lemma 1 For $r < n$ it is valid

$$e_r^2 X_{r+1} - (q + q^{-1}) e_r X_{r+1} e_r + X_{r+1} e_r^2 = 0.$$

Proof: For $r = n - 1$ it is true from the definition of (1). Now by an induction. We suppose it is valid for $r = k + 1$ and we are calculating

$$e_k^2 X_{k+1} - (q + q^{-1}) e_k X_{k+1} e_k + X_{k+1} e_k^2$$

from a definition X_{k+1} we obtain

$$\begin{aligned} &= e_k^2 (e_{k+1} X_{k+2} - q X_{k+2} e_{k+1}) - (q + q^{-1}) e_k (e_{k+1} X_{k+2} - q X_{k+2} e_{k+1}) e_k + \\ &\quad + (e_{k+1} X_{k+2} - q X_{k+2} e_{k+1}) e_k^2 \end{aligned}$$

from the (4) e_r commute with X_{r+2} and we have

$$\begin{aligned} &= [e_k^2 X_{k+1} - (q + q^{-1}) e_k X_{k+1} e_k + X_{k+1} e_k^2] X_{r+2} - \\ &\quad - q [e_k^2 X_{k+1} - (q + q^{-1}) e_k X_{k+1} e_k + X_{k+1} e_k^2] = 0 \end{aligned}$$

if we use the induction condition. ■

Simply from lemma 1 we obtain

$$e_r X_r = e_r^2 X_{r+1} - q e_r X_{r+1} e_r = q^{-1} (e_r X_{r+1} - q X_{r+1} e_r) e_r = q^{-1} X_r e_r \quad (5)$$

for $r < n$.

Similarly it is possible to prove that

$$e_r X_s = X_s e_r \text{ for } r + 1 > s. \quad (6)$$

We will continue the calculation of commutation between f_r and X_s .

From the definition (4) and the commutation relations (1) we have

$$f_r X_s = X_s f_r \text{ for } s > r. \quad (7)$$

If $r = s$ the calculation is more complicated than using the definition and the above relations gives

$$\begin{aligned} f_r X_r &= f_r (e_r X_{r+1} - q X_{r+1} e_r) = (e_r X_{r+1} - q X_{r+1} e_r) f_r - \\ &\quad \frac{-[(k_r - k_r^{-1}) X_{r+1} - q X_{r+1} (k_r - k_r^{-1})]}{(q - q^{-1})} = \\ &= X_r f_r - X_{r+1} \frac{(q^{-1} k_r - q k_r^{-1} - q k_r + q k_r^{-1})}{(q - q^{-1})} \end{aligned}$$

and finally we obtain

$$f_r X_r = X_r f_r + X_{r+1} k_r \quad (8)$$

In the last case $s < r$ e_s and f_r commute and if we use the definition of X_s we obtain

$$f_r X_s = X_s f_r + e_s [f_r, X_{s+1}] - q [f_r, X_{s+1}] e_s$$

If we put $s = r - 1$ we have

$$\begin{aligned} f_r X_{r-1} &= X_{r-1} f_r + e_{r-1} X_{r+1} k_r - q X_{r+1} k_r e_{r-1} = \\ X_{r-1} f_r + X_{r+1} (e_{r-1} k_r - q k_r e_{r-1}) &= X_{r-1} f_r \end{aligned}$$

By a simple induction we prove

$$f_r X_s = X_s f_r \text{ for } s < r \quad (9)$$

Now we have the all commutation relation which we need for the explicit construction of representations ϱ . The next lemma give the explicit form of the commutation relations e_i, f_i and k_i with $X_j^{m_j}$

Lemma 2 For $r < n$ it is valid

$$e_r X_r^{m_r} = q^{-m_r} X_r^{m_r} e_r,$$

$$e_r X_s^{m_s} = X_s^{m_s} e_r \text{ for } r \neq s-1, s,$$

$$e_r X_{r+1}^{m_{r+1}} = [m_{r+1}]_q X_{r+1}^{m_{r+1}-1} X_r + q^{m_{r+1}} X_{r+1}^{m_{r+1}} e_r,$$

$$f_r X_s^{m_s} = X_s^{m_s} f_r \text{ for } r \neq s, \quad (10)$$

$$f_r X_r^{m_r} = X_r^{m_r} f_r + [m_r]_q X_{r+1} X_r^{m_r-1} k_r,$$

$$k_r X_{r+1}^{m_{r+1}} = q^{-m_{r+1}} X_{r+1}^{m_{r+1}} k_r, k_r X_r^{m_r} = q^{m_r} X_r^{m_r} k_r,$$

$$k_r X_s^{m_s} = X_s^{m_s} k_r, \text{ for } s \neq r, r+1.$$

The special cases are f_n and k_n . In these cases we define

$$Y_{n-1} = e_{n-1}$$

and recurrently

$$Y_k = e_k Y_{k+1} - q Y_{k+1} e_k$$

for $k < n-1$ and it is valid

$$f_n X_n^{m_n} = X_n^{m_n} f_n - \frac{[m_n]_q}{(q - q^{-1})} X_n^{m_n-1} [q^{m_n-1} k_n - q^{-m_n+1} k_n^{-1}]$$

$$f_n X_r^{m_r} = X_r^{m_r} f_n - q^{-m_r+1} [m_r]_q X_r^{m_r-1} k_n^{-1} Y_r$$

$$k_n X_n^{m_n} = q^{2m_n} X_n^{m_n} k_n, k_n X_r^{m_r} = q^{m_r} X_r^{m_r} k_n,$$

Proof: By using the relations (4),(5),(6),(7),(8),(9) and an induction. The relations of k_r directly from the definition (1). ■

Using the relations (10) of lemma 1 we obtain the explicit form of the representation ϱ

$$\varrho(e_r)|m\rangle \otimes w = [m_{r+1}]_q |m - 1_{r+1} + 1_r\rangle \otimes w + q^{m_{r+1}-m_r} |m\rangle \otimes \phi(e_r)w,$$

$$\varrho(f_r)|m\rangle \otimes w = [m_r]_q |m + 1_{r+1} - 1_r\rangle \otimes \phi(k_r)w + |m\rangle \otimes \phi(f_r)w,$$

$$\varrho(k_r)|m\rangle \otimes w = q^{m_r-m_{r+1}} |m\rangle \otimes \phi(k_r)w,$$

$$\varrho(e_n)|m\rangle \otimes w = |m + 1_n\rangle \otimes w, \quad (11)$$

$$\begin{aligned}
\varrho(f_n)|m\rangle \otimes w &= - \frac{[m_n]_q}{(q - q^{-1})} |m - 1_n\rangle \otimes (q^{-1+\sum_{i=1}^n m_i} \phi(k_n) - q^{1-\sum_{i=1}^n m_i} \phi(k_n^{-1}))w - \\
&\quad - \sum_{k=1}^n [m_k]_q q^{1-\sum_{i=1}^k m_i} |m - 1_k\rangle \otimes \phi(k_{n+1}^{-1} Y_r)w, \\
\varrho(k_n)|m\rangle \otimes w &= q^{m_n+\sum_{i=1}^n m_i} |m\rangle \otimes \phi(k_n)w,
\end{aligned}$$

From the explicit form of the representation ϱ we can see that it is possible to rewrite this representation using the representation Γ (3). This representation Γ and representation ϕ are faithful representations and we can formulate the following theorem.

Theorem The mapping τ defined by formulae

$$\tau(e_r) = q^{N_{r+1}-N_r} e_r + b_r^+ b_{r+1}$$

$$\tau(f_r) = f_r + b_{r+1}^+ b_r k_r$$

$$\tau(k_r) = q^{N_r-N_{r+1}} K_r$$

for $r < n$

$$\tau(e_n) = b_n^+ \tag{12}$$

$$\tau(k_n) = q^{N_n+\sum_{i=1}^n N_i} k_n,$$

$$\tau(f_n) = - \frac{q^{\sum_{i=1}^n N_i} k_n - q^{-\sum_{i=1}^n N_i} k_n^{-1}}{(q - q^{-1})} b_n - \sum_{k=1}^{n-1} q^{-\sum_{i=1}^k N_i} k_n^{-1} Y_k b_k$$

is a homomorphism from $U_q(sl(n+1, C))$ to $W_{2(n)}^q \otimes U_q(gl(n, C))$.

In this letter we have presented some simple generalization of the construction (Fu and Ge 1992). The realizations of $U_q(sl(n+1, C))$ are in q-boson pairs and in the generators of the subalgebra $U_q(gl(n, C))$. For using our formulae (12) recurrently to obtain the pure q-boson realizations it is also needed to have an operator $\tau(k_{n+1})$. Evidently from our construction it is possible reformulate for $U_q(gl(n+1, C))$ and after calculation we obtain

$$\tau(k_{n+1}) = q^{\lambda_{n+1}-\sum_{i=1}^n N_i}$$

The pure q-boson realizations are a starting point (see Fu and Ge 1992) for a construction of the cyclic representations in the root of unity. The properties of these representations we will study in a forthcoming paper.

The authors are grateful to the members of the seminar of the quantum groups in Prague especially Dr M.Havlíček ,for useful discussions.

References

- Biedernharn L C 1989** J.Phys.A:Math.Gen. 22 p.L873
- Burdík Ć Havlíček M and Vančura T 1992** Commun.Math.Phys. in print
- Burdík Ć and Navrátil O 1990** J.Phys.A:Math.Gen. 23 pL1205
- Burdík Ć 1985** J.Phys.A:Math.Gen. 18 p.3101
- Burroughs N 1990** Commun.Math.Phys.127 p.109
- Hayashi T 1990** Commun.Math.Phys.127 p.129
- Hong-Chen Fu and Mo-Lin Ge 1992** J.Math.Phys. 33 p.427
- Rosso M 1988** Commun.Math.Phys.117 p.581
- Rosso M 1989** Commun.Math.Phys.124 p.307